

**PAIN MEDICATION  
Diluted Meloxicam\***

**\*\*10/14/2023\*\***

After cat is under anesthesia, and prior to surgery prep, weigh cat and administer diluted meloxicam. Administer SubQ.

**△ Caution: do not administer to sick or dehydrated cats. If cat is older and may have kidney disease, consult prior to administering.**

Choose lower dosage if weight falls in between lbs.

<b>Wt in lbs</b>	<b>cc/ml</b>
<b>2</b>	<b>.04</b>
<b>4</b>	<b>.25</b>
<b>6</b>	<b>.40</b>
<b>8</b>	<b>.45</b>
<b>10</b>	<b>.70</b>
<b>12</b>	<b>.80</b>
<b>14</b>	<b>.85</b>
<b>16</b>	<b>1.10</b>
<b>18</b>	<b>1.25</b>
<b>20</b>	<b>1.35</b>

\*Diluted 5:1 from stock solution using 12 cc syringe.  
To make: draw up 2 cc, then 8 ml sterile water.  
Put in orange bot

**PAIN MEDICATION  
Diluted Meloxicam\***

**\*\*10/14/2023\*\***

After cat is under anesthesia, and prior to surgery prep, weigh cat and administer diluted meloxicam. Administer SubQ.

**△ Caution: do not administer to sick or dehydrated cats. If cat is older and may have kidney disease, consult prior to administering.**

Choose lower dosage if weight falls in between lbs.

<b>Wt in lbs</b>	<b>cc/ml</b>
<b>2</b>	<b>.04</b>
<b>4</b>	<b>.25</b>
<b>6</b>	<b>.40</b>
<b>8</b>	<b>.45</b>
<b>10</b>	<b>.70</b>
<b>12</b>	<b>.80</b>
<b>14</b>	<b>.85</b>
<b>16</b>	<b>1.10</b>
<b>18</b>	<b>1.25</b>
<b>20</b>	<b>1.35</b>

\*Diluted 5:1 from stock solution using 12 cc syringe.  
To make: draw up 2 cc, then 8 ml sterile water.  
Put in orange bottle.